

Allergy and A	naphylaxis Emergency Plan	BEDFORD CITY SCHOOL DISTRICT
School:	Grade/Room:	PROUDLY SERVING BEDFORD + BEDFORD HTS. • WALTON HILLS • OAKWOO
Child's name:	Date of	plan:

Child's name:	Date of plan:	
Date of birth:/ Ag	je Weight:	Attach child's
Child has allergy to		photo
Child has asthma. Child has had anaphylaxis. Child may carry medicine.	☐ Yes ☐ No (If yes, higher chance severe reaction) ☐ Yes ☐ No ☐ Yes ☐ No	
	☐ Yes ☐ No (If child refuses/is unable to self-treat, an adult n	nust give medicine)

#### IMPORTANT REMINDER

Anaphylaxis is a potentially life-threating, severe allergic reaction. If in doubt, give epinephrine.

#### For Severe Allergy and Anaphylaxis What to look for



If child has ANY of these severe symptoms after eating the food or having a sting, give epinephrine.

- Shortness of breath, wheezing, or coughing
- Skin color is pale or has a bluish color
- Weak pulse
- Fainting or dizziness
- Tight or hoarse throat
- Trouble breathing or swallowing
- Swelling of lips or tongue that bother breathing
- Vomiting or diarrhea (if severe or combined with other symptoms)
- Many hives or redness over body
- Feeling of "doom," confusion, altered consciousness, or agitation

☐ **SPECIAL SITUATION**: If this box is checked, child has an extremely severe allergy to an insect sting or the following food(s): . Even if child has MILD symptoms after a sting or eating these foods, give epinephrine.

#### Give epinephrine! What to do

- 1. Inject epinephrine right away! Note time when epinephrine was given.
- 2. Call 911.
  - Ask for ambulance with epinephrine.
  - Tell rescue squad when epinephrine was given.
- 3. Stav with child and:
  - Call parents and child's doctor.
  - Give a second dose of epinephrine, if symptoms get worse, continue, or do not get better in 5 minutes.
  - Keep child lying on back. If the child vomits or has trouble breathing, keep child lying on his or her side.
- 4. Give other medicine, if prescribed. Do not use other medicine in place of epinephrine.
  - Antihistamine
  - Inhaler/bronchodilator

#### For Mild Allergic Reaction What to look for



If child has had any mild symptoms, monitor child. Symptoms may include:

- Itchy nose, sneezing, itchy mouth
- A few hives
- Mild stomach nausea or discomfort

### Monitor child What to do

Stay with child and:

- Watch child closely.
- Give antihistamine (if prescribed).
- Call parents and child's doctor.
- If more than 1 symptom or symptoms of severe allergy/anaphylaxis develop, use epinephrine. (See "For Severe Allergy and Anaphylaxis.")

Parent/Guardian Authorization Signature	Date	Physician/HCP Authorization Signature	 Date
Other (for example, inhaler/bronchodilator if child has ast	thma):		
Antihistamine, by mouth (type and dose):		(*Use 0.15 mg, if 0.10 mg is not ava	ailable)
		$\square$ 0.30 mg (25 kg or more)	
		$\square$ 0.15 mg (13 kg to less than 2	25 kg)
Epinephrine, intramuscular (list type):		Dose: $\square$ 0.10 mg (7.5 kg to less than	13 kg)*
Medicines/Doses			

Parent/Guardian Authorization Signature

Physician/HCP Authorization Signature

# Allergy and Anaphylaxis Emergency Plan



Child's name:	Date of plan:	
Additional Instructions:		
Contacts		
Call 911 / Rescue squad:		
Healthcare Provider:	Phone:	
Parent/Guardian:	Phone:	
Parent/Guardian:	Phone:	
Other Emergency Contacts		
Name/Relationship:	Phone:	
Name/Relationship:	Phone:	

If the student has been instructed on how to use the Epinephrine Auto-Injector and is able to self-adminster, please complete the following page. This is required for your student to self-carry an Epinephrine Auto-Injector at school or during a school event. If the student is able to self carry, it is required by law that an additional Epinephrine auto injector is kept in the school health clinic.



## Authorization for Student Possession and Use of an Epinephrine Autoinjector

In accordance with ORC 3313.718/3313.141

Student name	
Student address	
This section must be completed and signed by the ${f s}$	student's parent or guardian.
at the school and any activity, event, or program sponso	whild to possess and use an epinephrine autoinjector, as prescribed, bred by or in which the student's school is a participant. I understand ance from an emergency medical service provider if this medication dication to the school nurse as required by law.
Parent / Guardian signature	Date
Parent/Guardian name	Parent/Guardian emergency telephone number
	( )
This section must be completed and signed by the n	nedication prescriber.
Name and dosage of medication	
Date medication administration begins	Date medication administration ends (if known)
Date medication administration begins	Date medication administration ends (if known)
Circumstances for use of the epinephrine autoinjector	1
Procedures for school employees if the student is unable to administer	
Procedures for school employees if the student is unable to administer	the medication or init does not produce the expected relief
3	
Possible severe adverse reactions:	
To the student for which it is prescribed (that should be reported to the	prescriber)
To a student for which it is <b>not</b> prescribed who receives a dose	
Special instructions	
As the prescriber, I have determined that this student and have provided the student with training in the provided that this student with the stu	t is capable of possessing and using this autoinjector appropriately
Prescriber signature	Date
Prescriber name	Prescriber emergency telephone number